



Using leverages for objective analysis of PSMSL tide gauges in Arctic Ocean sea level reconstruction

Svendsen, Peter Limkilde; Andersen, Ole Baltazar; Nielsen, Allan Aasbjerg

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Svendsen, P. L., Andersen, O. B., & Nielsen, A. A. (2013). *Using leverages for objective analysis of PSMSL tide gauges in Arctic Ocean sea level reconstruction*. Poster session presented at PSMSL 80th Anniversary Workshop on Sea Level Science, Liverpool, United Kingdom.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Using leverages for objective analysis of PSMSL tide gauges in Arctic Ocean sea level reconstruction

Technical University
of Denmark



Peter Limkilde Svendsen¹, Ole Baltazar Andersen¹ and Allan Aasbjerg Nielsen²
plsv@space.dtu.dk, oa@space.dtu.dk, alan@dtu.dk | ¹DTU Space, ²DTU Compute
Technical University of Denmark, Kgs. Lyngby, Denmark

INTRODUCTION


For reconstructing historical sea levels in the Arctic area, lack of data presents a major challenge. We attempt to adapt the model by Church et al. (2004), examining inclusion criteria for tide gauges in the area. The reconstruction model is based upon spatial, stationary patterns of variability extracted from a calibration period, usually satellite data. These patterns are determined as empirical orthogonal functions (EOFs). The model determines, for each point in time, an appropriately weighted sum of these, constrained locally by tide gauge records and regularized per Kaplan et al. (1997). The *leverage* of each tide gauge is a statistical measure of its influence on the result. This way, we can readily identify possible outliers among the tide gauge records in a procedural, objective way.

DATA


We use monthly PSMSL tide gauge data (RLR only) above approx. 60°N, allowing only records from stations with more than 5 years of data. We use the first differences of the time series to work around vertical datum considerations, then integrate the results, as in Church et al. (2004). GIA correction has been applied to the tide gauges, using the Peltier ICE-5G model (Peltier, 2004). For this preliminary analysis, the EOFs have been extracted from the Drakkar ocean model (1949–2008).

EOFs

The reconstruction model incorporates a spatially uniform pattern (known as an “EOF0”), in addition to the leading ten EOFs. The first four of these are shown below.



Time series of EOFs




EOF1 is fairly close to uniform and may capture a large amount of overall trend. It is by far the dominant mode in the Drakkar data, explaining almost 90% of the variance.

HIGH-LEVERAGE GAUGES

In least-squares regression, the leverage of each observation is given by the diagonal elements of the “hat matrix” that relates the response variable y (in this case, tide gauge records) to its estimate, \hat{y} . See e.g. Nielsen, 2013.


The map shows all tide gauges included in the reconstruction (green) and those rejected due to high leverage (red). Rejection criterion here is three times the mean gauge leverage.

The plot shows an example of rejected gauges, the three on Kotelný Island (75°N, 140°E). These gauges may simply not be sufficiently represented within the appropriate EOFs.



LOCAL RECONSTRUCTION ABILITY

To assess the validity of the results, the reconstructed sea-level time series in selected points have been plotted and compared with actual tide gauge data and the existing MONARCH-A reconstruction (by Henry et al.). The model has been applied both without regularization (“OLS”) and with regularization (“Kaplan”).



The unregularized reconstruction seems more prone to developing nonexistent trends in the time series, and not very appropriate for this purpose. Removing a handful of gauges of high leverage seems to dampen the oscillations near the coast considerably, while keeping almost the same behaviour in open ocean.

REFERENCES

- Church, J., White, N., Coleman, R., Lambeck, K. and Mitrović, J. (2004). Estimates of the regional distribution of sea level rise over the 1950–2000 period. *Journal of Climate*, Volume 17, Issue 13, pp. 2609–2625.
- Kaplan, A., Kushni, Y., Cane, M.A. and Blumenthal, M.B. (1997). Reduced space optimal analysis for historical data sets: 136 years of Atlantic sea surface temperatures. *Journal of Geophysical Research*, Volume 102, Issue C13, pp. 27,835–27,860.
- Nielsen, A.A. (2013). Least Squares Adjustment: Linear and Nonlinear Weighted Regression Analysis. Lecture note.
- Peltier, W.R. (2004). Global glacial isostasy and the surface of the ice-age Earth: The ICE-5G (VM2) model and GRACE. *Annual Review of Earth and Planetary Sciences*, Volume 32, pp. 111–149.